

Activity of *Cissus quadrangularis* against Herpes through deactivation of Thymidine Kinase of Herpes Simplex virus (1KIM)

Chinmayee Chitralkha Pradhan¹, Debanjana Saha²

¹pradhanchinmayee96@gmail.com

²deepanjana.saha@cutm.ac.in

Centurion University of Technology and Management, Odisha, India

Abstract: An in-silico study was performed to determine the activity of *Cissus quadrangularis* against Herpes. Molecular docking using Biovia Discovery Studio was performed to identify the phytochemical responsible to deactivate Thymidine Kinase of Herpes Simplex virus (1KIM) enzyme. It was found that Ferulic acid helped to prevent Herpes.

Introduction: *Cissus quadrangularis* is known for its medicinal activities. The roots and stems are most useful for healing of fracture of the bones. The stem is bitter; it is given internally and applied topically in broken bones, used in complaints of the back and spine. A paste of stem is useful for muscular pains and herpes. The plant has been documented in Ayurveda for the treatment of osteoarthritis, rheumatoid arthritis and osteoporosis.

The plant is classified as follows:

Kingdom	Plantae
Division	Tracheophyta
Class	Magnoliopsida
Order	Vitales
Family	Vitaceae
Genus	<i>Cissus</i>
Species	<i>quadrangularis</i>

Major phytochemicals present in the plant are:

- a. Lupeol
- b. Ferulic acid
- c. Hesperidin
- d. Naringin

One of the major enzymes required for the survival of the organism causing Herpes is Thymidine Kinase of Herpes Simplex virus (1KIM) enzyme. The objective of this work is to find the phytochemical that can deactivate the enzyme, thereby preventing the physiological activity of the organism.