Activity of Salvia officinalis against Herpes through deactivation of Thymidine Kinase of Herpes Simplex virus (1KIM)

Prasityaditya Mohanty¹, Pratap Kumar Chhotaray²

¹prasityadityamohanty1997@gmail.com

²pratap.chottaray@cutm.ac.in

Centurion University of Technology and Management, Odisha, India

Abstract: An in-silico study was performed to determine the activity of Salvia officinalis against Herpes. Molecular docking using Biovia Discovery Studio was performed to identify the phytochemical responsible to deactivate Thymidine Kinase of Herpes Simplex virus (1KIM) enzyme. It was found that Tocopherol helped to prevent Herpes.

Introduction: Salvia officinalis is known for its medicinal activities. S. officinalis has been used for the treatment of different kinds of disorders including seizure, ulcers, gout, rheumatism, herpes, inflammation, dizziness, tremor, paralysis, diarrhea, and hyperglycemia.

The plant is classified as follows:

Kingdom	Plantae
Division	Tracheophyta
Class	Magnoliopsida
Order	Lamiales
Family	Lamiaceae
Genus	Salvia
Species	officinalis

Major phytochemicals present in the plant are:

- a. Tocopherol
- b. Epicatechin
- c. Coumarin
- d. Proanthocyanidins

One of the major enzymes required for the survival of the organism causing Herpes is Thymidine Kinase of Herpes Simplex virus (1KIM) enzyme. The objective of this work is to find the phytochemical that can deactivate the enzyme, thereby preventing the physiological activity of the organism.

ISSN: 2395-6216