Activity of Hottuynia cordata against COVID 19 through deactivation of papain-like protease of SARS CoV-2 (6W9C)

Aditya Narayan Barik¹, Yashaswi Nayak²

¹190705180086@cutm.ac.in

²yashaswi.nayak@cutm.ac.in

Centurion University of Technology and Management, Odisha, India

Abstract: An in-silico study was performed to determine the activity of Hottuynia cordata against COVID 19. Molecular docking using Biovia Discovery Studio was performed to identify the phytochemical responsible to deactivate papain-like protease of SARS CoV-2 (6W9C) enzyme. It was found that Coumarin helped to prevent COVID 19.

Introduction: Hottuynia cordata is known for its medicinal activities. It is used as a fresh herbal garnish. In northeastern India, it is commonly used in salads and as a garnish over side dishes. The tender roots can also be ground into chutneys along with dry meat or fish, chilies, and tamarind. It is taken raw as salad and cooked along with fish as fish curry. In Japan and Korea, its dried leaves may be used as a tea. Houttuynia cordata was used in traditional Chinese medicine.

The plant is classified as follows:

Kingdom	Plantae
Division	Tracheophytes
Class	Angiosperms
Order	Piperales
Family	Saururaceae
Genus	Houttuynia
Species	cordata

Major phytochemicals present in the plant are:

- a. Lupeol
- b. Peonidin
- c. Coumarin
- d. Malvidin

One of the major enzymes required for the survival of the organism causing COVID 19 is papainlike protease of SARS CoV-2 (6W9C) enzyme. The objective of this work is to find the phytochemical that can deactivate the enzyme, thereby preventing the physiological activity of the organism.

> Centurion Journal of Multidisciplinary Research Special Issue: June 2020

ISSN: 2395-6216